

**ab226766 –
PD-L1 RabMAb[®]
antibody (clone 73-10)**

Protocol Booklet

For Immunohistochemistry.

This product is for research use only and is not intended for diagnostic use.

Ventana Benchmark XT

1. Perform heat mediated antigen retrieval in sCC1 (Tris/EDTA buffer, pH 8).
2. Use ab226766 at a dilution of 10µg/ml.
3. Incubate for 30 minutes at 37°C.
4. Signal detection with BenchMark XT from Roche/Ventana and ultraView Universal DAB Detection Kit (Product code: 760-500)

Dako Autostainer Link 48

1. Perform heat mediated antigen retrieval in high pH buffer (Tris/EDTA buffer, pH 9, during 20 min at 95°C).
2. Use ab226766 at a dilution of 2µg/ml.
3. Incubate for 30 minutes at room temperature.
4. Block sample with peroxidase blocking buffer (EnVision Flex Peroxidase-Blocking Reagent) for 5 minutes.
5. Signal detection with Autostainer Link from Dako and EnVision Flex Kit, High pH (Product code: K8000).

UK, EU and ROW

Email: technical@abcam.com

Tel: +44 (0)1223 696000

www.abcam.com

US, Canada and Latin America

Email: us.technical@abcam.com

Tel: 888-77-ABCAM (22226)

www.abcam.com

China and Asia Pacific

Email: hk.technical@abcam.com

Tel: 108008523689 (中國聯通)

www.abcam.cn

Japan

Email: technical@abcam.co.jp

Tel: +81-(0)3-6231-0940

www.abcam.co.jp